I. Multiple Choices (40%)

1. If X and Y are any random variables with $E(X) = 5$, $E(Y) = 6$, $E(XY) = 21$, $V(X) = 9$ and $V(Y) = 10$, then the relationship between X and Y is a:
   (A) strong positive relationship.
   (B) strong negative relationship.
   (C) weak positive relationship.
   (D) weak negative relationship.

2. When extreme values are present in a set of data, which of the following descriptive summary measures are most appropriate?
   (A) Coefficient of variation and range
   (B) Mean and standard deviation
   (C) Interquartile range and median
   (D) Variance and interquartile range

3. If the standard error of estimate $s_e = 20$ and $n = 10$, then the sum of squares for error, SSE, is:
   (A) 400
   (B) 3,200
   (C) 4,000
   (D) 40,000

4. The standardized residual is defined as:
   (A) residual divided by the standard error of estimate.
   (B) residual multiplied by the square root of the standard error of estimate.
   (C) residual divided by the square of the standard error of estimate.
   (D) residual multiplied by the standard error of estimate.

5. The model $y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \beta_3 x_1 x_2 + \varepsilon$　is referred to as a:
   (A) first-order model with two predictor variables with no interaction.
   (B) first-order model with two predictor variables with interaction.
   (C) second-order model with three predictor variables with no interaction.
   (D) second-order model with three predictor variables with interaction.

6. The sign test statistic is approximately normally distributed whenever the sample sizes are larger than:
   (A) 5
   (B) 10
   (C) 30
   (D) None of these choices.

7. Which of the following tests employs matched-pairs sampling?
   (A) The Wilcoxon signed-rank sum test
   (B) The Wilcoxon rank-sum test
   (C) The Mann-Whitney test
   (D) None of these choices

8. When is the Tukey multiple comparison method used?
   (A) To test for normality.
   (B) To test equal variances.
   (C) To test for equality of a group of population means.
   (D) To test for differences in pairwise means.

9. Which statistical technique is appropriate when we wish to analyze the relationship between two qualitative variables with two or more categories?
   (A) The chi-squared test of a multinomial experiment
   (B) The chi-squared test of a contingency table
   (C) The t-test of the difference between two means
   (D) The z-test of the difference between two proportions

10. Fisher's least significant difference (LSD) multiple comparison method is flawed because
    (A) it will increase α, the probability of committing a Type I error.
    (B) it will increase β, the probability of committing a Type II error.
    (C) it will increase both α and β, the probabilities of committing Type I and Type II errors, respectively.
    (D) None of these choices.

II. Short Answers (60%)

1. Test the hypotheses whether there is no first-order autocorrelation in regression residuals, given that: Durbin-Watson Statistic d = 1.12, n = 45, k = 5, and α = 0.05. (10%)

2. Three people toss a coin and the odd man pays for coffee. If the coins all turn up the same, they are tossed again. Find the probability that fewer than 4 tosses are needed. (10%)

3. In a random sample of n=500 families owning television sets in the city of Hamilton, Canada, it is found that x=340 subscribed to HBO. (10%)
   (a) Find a 95% confidence interval for the actual proportion of families in this city who subscribe to HBO.
   (b) How large a sample is required if we want to be 95% confident that our estimate of p is within 0.02?

4. A partial ANOVA table in a randomized block design is shown below, where the treatments refer to different high blood pressure drugs, and the blocks refer to groups of men with similar levels of high blood pressure. (20%)

Source of Variation	SS	df	MS	F
Treatments	*	4	*	*
Blocks	3,120	6	*	*
Error	*	*	115	
Total	12,600	34		

   (a) Fill in the missing values (identified by asterisks) in the above ANOVA Table.
   (b) Can we infer at the 5% significance level that the treatment means differ?
   (c) Can we infer at the 5% significance level that the block means differ?

5. Suppose the probability is 0.8 that any given person will believe a tale about the transgression of a famous actress. What is the probability that (10%)
   (a) The sixth person to hear this tale is the fourth one to believe it?
   (b) The third person to hear this tale is the first one to believe it?

Durbin-Watson Statistic: 5 Per Cent Significance Points of dL and dU

	k'*=1		k'=2		k'=3		k'=4		k'=5		k'=6		k'=7		k'=8	
n	dL	dU	dL	dU	dL	dU	dL	dU	dL	dU	dL	dU	dL	dU	dL	dU
6	0.610	1.400	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
7	0.700	1.356	0.467	1.896	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
8	0.763	1.332	0.559	1.777	0.367	2.287	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
9	0.824	1.320	0.629	1.699	0.455	2.128	0.296	2.588	-----	-----	-----	-----	-----	-----	-----	-----
10	0.879	1.320	0.697	1.641	0.525	2.016	0.376	2.414	0.243	2.822	-----	-----	-----	-----	-----	-----
11	0.927	1.324	0.758	1.604	0.595	1.928	0.444	2.283	0.315	2.645	0.203	3.004	-----	-----	-----	-----
12	0.971	1.331	0.812	1.579	0.658	1.864	0.512	2.177	0.380	2.506	0.268	2.832	0.171	3.149	-----	-----
13	1.010	1.340	0.861	1.562	0.715	1.816	0.574	2.094	0.444	2.390	0.328	2.692	0.230	2.985	0.147	3.266
14	1.045	1.350	0.905	1.551	0.767	1.779	0.632	2.030	0.505	2.296	0.389	2.572	0.286	2.848	0.200	3.111
15	1.077	1.361	0.946	1.543	0.814	1.750	0.685	1.977	0.562	2.220	0.447	2.471	0.343	2.727	0.251	2.979
16	1.106	1.371	0.982	1.539	0.857	1.728	0.734	1.935	0.615	2.157	0.502	2.388	0.398	2.624	0.304	2.860
17	1.133	1.381	1.015	1.536	0.897	1.710	0.779	1.900	0.664	2.104	0.554	2.318	0.451	2.537	0.356	2.757
18	1.158	1.391	1.046	1.535	0.933	1.696	0.820	1.872	0.710	2.060	0.603	2.258	0.502	2.461	0.407	2.668
19	1.180	1.401	1.074	1.536	0.967	1.685	0.859	1.848	0.752	2.023	0.649	2.206	0.549	2.396	0.456	2.589
20	1.201	1.411	1.100	1.537	0.998	1.676	0.894	1.828	0.792	1.991	0.691	2.162	0.595	2.339	0.502	2.521
21	1.221	1.420	1.125	1.538	1.026	1.669	0.927	1.812	0.829	1.964	0.731	2.124	0.637	2.290	0.546	2.461
22	1.239	1.429	1.147	1.541	1.053	1.664	0.958	1.797	0.863	1.940	0.769	2.090	0.677	2.246	0.588	2.407
23	1.257	1.437	1.168	1.543	1.078	1.660	0.986	1.785	0.895	1.920	0.804	2.061	0.715	2.208	0.628	2.360
24	1.273	1.446	1.188	1.546	1.101	1.656	1.013	1.775	0.925	1.902	0.837	2.035	0.750	2.174	0.666	2.318
25	1.288	1.454	1.206	1.550	1.123	1.654	1.038	1.767	0.953	1.886	0.868	2.013	0.784	2.144	0.702	2.280
26	1.302	1.461	1.224	1.553	1.143	1.652	1.062	1.759	0.979	1.873	0.897	1.992	0.816	2.117	0.735	2.246
27	1.316	1.469	1.240	1.556	1.162	1.651	1.084	1.753	1.004	1.861	0.925	1.974	0.845	2.093	0.767	2.216
28	1.328	1.476	1.255	1.560	1.181	1.650	1.104	1.747	1.028	1.850	0.951	1.959	0.874	2.071	0.798	2.188
29	1.341	1.483	1.270	1.563	1.198	1.650	1.124	1.743	1.050	1.841	0.975	1.944	0.900	2.052	0.826	2.164
30	1.352	1.489	1.284	1.567	1.214	1.650	1.143	1.739	1.071	1.833	0.998	1.931	0.926	2.034	0.854	2.141
31	1.363	1.496	1.297	1.570	1.229	1.650	1.160	1.735	1.090	1.825	1.020	1.920	0.950	2.018	0.879	2.120
32	1.373	1.502	1.309	1.574	1.244	1.650	1.177	1.732	1.109	1.819	1.041	1.909	0.972	2.004	0.904	2.102
33	1.383	1.508	1.321	1.577	1.258	1.651	1.193	1.730	1.127	1.813	1.061	1.900	0.994	1.991	0.927	2.085
34	1.393	1.514	1.333	1.580	1.271	1.652	1.208	1.728	1.144	1.808	1.079	1.891	1.015	1.978	0.950	2.069
35	1.402	1.519	1.343	1.584	1.283	1.653	1.222	1.726	1.160	1.803	1.097	1.884	1.034	1.967	0.971	2.054
36	1.411	1.525	1.354	1.587	1.295	1.654	1.236	1.724	1.175	1.799	1.114	1.876	1.053	1.957	0.991	2.041
37	1.419	1.530	1.364	1.590	1.307	1.655	1.249	1.723	1.190	1.795	1.131	1.870	1.071	1.948	1.011	2.029
38	1.427	1.535	1.373	1.594	1.318	1.656	1.261	1.722	1.204	1.792	1.146	1.864	1.088	1.939	1.029	2.017
39	1.435	1.540	1.382	1.597	1.328	1.658	1.273	1.722	1.218	1.789	1.161	1.859	1.104	1.932	1.047	2.007
40	1.442	1.544	1.391	1.600	1.338	1.659	1.285	1.721	1.230	1.786	1.175	1.854	1.120	1.924	1.064	1.997
45	1.475	1.566	1.430	1.615	1.383	1.666	1.336	1.720	1.287	1.776	1.238	1.835	1.189	1.895	1.139	1.958
50	1.503	1.585	1.462	1.628	1.421	1.674	1.378	1.721	1.335	1.771	1.291	1.822	1.246	1.875	1.201	1.930

F distribution Table

F Values for $\alpha = 0.05$

d_2	1	2	3	4	5	6	7	8	9
1	161.4	199.5	215.7	224.6	230.2	234.0	236.8	238.9	240.5
2	18.51	19.00	19.16	19.25	19.3	19.33	19.35	19.37	19.38
3	10.13	9.55	9.28	9.12	9.01	8.94	8.89	8.85	8.81
4	7.71	6.94	6.59	6.39	6.26	6.16	6.09	6.04	6.00
5	6.61	5.79	5.41	5.19	5.05	4.95	4.88	4.82	4.77
6	5.99	5.14	4.76	4.53	4.39	4.28	4.21	4.15	4.10
7	5.59	4.74	4.35	4.12	3.97	3.87	3.79	3.73	3.68
8	5.32	4.46	4.07	3.84	3.69	3.58	3.50	3.44	3.39
9	5.12	4.26	3.86	3.63	3.48	3.37	3.29	3.23	3.18
10	4.96	4.10	3.71	3.48	3.33	3.22	3.14	3.07	3.02
11	4.84	3.98	3.59	3.36	3.20	3.09	3.01	2.95	2.90
12	4.75	3.89	3.49	3.26	3.11	3.00	2.91	2.85	2.80
13	4.67	3.81	3.41	3.18	3.03	2.92	2.83	2.77	2.71
14	4.60	3.74	3.34	3.11	2.96	2.85	2.76	2.70	2.65
15	4.54	3.68	3.29	3.06	2.90	2.79	2.71	2.64	2.59
16	4.49	3.63	3.24	3.01	2.85	2.74	2.66	2.59	2.54
17	4.45	3.59	3.20	2.96	2.81	2.70	2.61	2.55	2.49
18	4.41	3.55	3.16	2.93	2.77	2.66	2.58	2.51	2.46
19	4.38	3.52	3.13	2.90	2.74	2.63	2.54	2.48	2.42
20	4.35	3.49	3.10	2.87	2.71	2.60	2.51	2.45	2.39
21	4.32	3.47	3.07	2.84	2.68	2.57	2.49	2.42	2.37
22	4.30	3.44	3.05	2.82	2.66	2.55	2.46	2.40	2.34
23	4.28	3.42	3.03	2.80	2.64	2.53	2.44	2.37	2.32
24	4.26	3.40	3.01	2.78	2.62	2.51	2.42	2.36	2.30
25	4.24	3.39	2.99	2.76	2.60	2.49	2.40	2.34	2.28
26	4.23	3.37	2.98	2.74	2.59	2.47	2.39	2.32	2.27
27	4.21	3.35	2.96	2.73	2.57	2.46	2.37	2.31	2.25
28	4.20	3.34	2.95	2.71	2.56	2.45	2.36	2.29	2.24
29	4.18	3.33	2.93	2.70	2.55	2.43	2.35	2.28	2.22
30	4.17	3.32	2.92	2.69	2.53	2.42	2.33	2.27	2.21
40	4.08	3.23	2.84	2.61	2.45	2.34	2.25	2.18	2.12
60	4.00	3.15	2.76	2.53	2.37	2.25	2.17	2.10	2.04
120	3.92	3.07	2.68	2.45	2.29	2.17	2.09	2.02	1.96
inf	3.84	3.00	2.60	2.37	2.21	2.10	2.01	1.94	1.88