國立彰化師範大學 97 學年度碩士班招生考試試題

☆☆請在答案紙上作答☆☆

共2頁,第1頁

1. Solve the following three sub-questions:

- (a) By using the technique of integration, find the volume of a right circular cone with base radius R and height H. (10%)
- (b) By employing the extreme-value theorem, find the volume of a right circular cylinder of greatest volume that can be inserted in the right circular cone as you derived in (a). (10%)
- (c) A display screen is to be laid out in a rectangular area and protected by a magnetic traction free wire frame. What is the largest possible area of the display screen if only 500cm of the wire is available for the frame? (10%)

Note: Please draw the corresponding diagrams to show your assumptions for the above sub-questions.

2. The trigonometric function f(t) is given by

$$f(t) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi t}{p} + b_n \sin \frac{n\pi t}{p} \right),$$

where a_0, a_n , and b_n are the Euler coefficients of f(t); 2p is the period of the function:

- (a) Show that the function f(t) can be written as $f(t) = A_0 + \sum_{n=1}^{\infty} A_n \sin(\frac{n\pi t}{p} + \theta_n)$. Please express the new defined coefficients, A_0 , A_n , θ_n in terms of a_0 , a_n , b_n . (15%)
- (b) Show that the function f(t) can be written as $f(t) = \sum_{n=-\infty}^{\infty} C_n e^{i\frac{n\pi}{p}}$, $i = \sqrt{-1}$, e = the natural exponential base. Please express the new defined coefficients, C_n , in terms of a_0, a_n, b_n . (15%)
- 3. The following three sub-problems are related to the Laplace transformation. It is recalled that the Laplace transform of a function f(t) associates a function of s is given by

$$L[f(t)] = F(s) = \int_{0}^{\infty} f(t)e^{-st}dt$$

(a) Find the Laplace transform of f(t) = u(t - a), where $u(t - a) = \begin{cases} 1, & t \ge a \\ 0, & t < a \end{cases}$. (10%)

國立彰化師範大學 97 學年度碩士班招生考試試題

系所: 顯示技術研究所碩士班

科目: 工程數學

☆☆請在答案紙上作答☆☆

共2頁,第2頁

- (b) Find the inverse transform of $F(s) = \frac{e^{-s}}{s^2 + 3s + 2}$. (10%)
- (c) By employing Laplace transformation, find the solution of the following differential equation which satisfies the given initial conditions,

$$y'' + 3y' + 2y = u(t-1)$$
, $y = y(t)$ and $y' = \frac{dy}{dt}$

with the given conditions y(0) = 0 and y'(0) = 1. (20%)